arXiv:gr-qc/0509012v1 2 Sep 2005

Parametric derivation of the observable relativistic periastron advance for binary pulsars

Christian Königsdörffer* and Achamveedu Gopakumar†
Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, Max-Wien-Platz 1, 07743 Jena, Germany
(Dated: 7th February 2008)

We compute the dimensionless relativistic periastron advance parameter k, which is measurable from the timing of relativistic binary pulsars. We employ for the computation the recently derived Keplerian-type parametric solution to the post-Newtonian (PN) accurate conservative dynamics of spinning compact binaries moving in eccentric orbits. The parametric solution and hence the parameter k are applicable for the cases of *simple precession*, namely, case (i), the binary consists of equal mass compact objects, having two arbitrary spins, and case (ii), the binary consists of compact objects of arbitrary mass, where only one of them is spinning with an arbitrary spin. Our expression, for the cases considered, is in agreement with a more general formula for the 2PN accurate k, relevant for the relativistic double pulsar PSR J0737–3039, derived by Damour and Schäfer many years ago, using a different procedure.

PACS numbers: 04.25.Nx, 97.60.Gb, 97.60.Jd, 97.80.-d

I. INTRODUCTION

A detailed knowledge about the dynamics of spinning compact binaries is of interest to both astrophysicists and theoretical physicists. The astrophysical interest arises from the highly accurate radio observations of relativistic binary pulsars, which can be used to test general relativity in strong field regimes [1]. Moreover, inspiralling spinning compact binaries are promising sources of gravitational radiation for the ground based and proposed space based laser interferometric gravitational-wave detectors [2]. The theoretical interest is due to the possibility of constructing numerically spinning compact binary configurations in full general relativity [see Ref. [3] and references therein]. Further, it is of some interest to explore if the dynamics of spinning compact binaries will be deterministic or not [4, 5].

In many of the above mentioned scenarios, the dynamics of spinning compact binaries can be described in great detail by the post-Newtonian (PN) approximation to general relativity. The PN approximation allows one to express the equations of motion of a compact binary as corrections to Newtonian equations of motion in powers of $(v/c)^2 \sim GM/(c^2R)$, where v, M, and R are the characteristic orbital velocity, the total mass and the typical orbital separation of the binary. For spinning compact binaries, the dynamics is determined, not only by the orbital equations of motion for these objects, but also by the precessional equations for the orbital plane and the spin vectors themselves [6]. Note that for spinning compact binaries, spin effects enter the dynamics for the first time through spin-orbit interaction. Therefore, the Hamiltonian describing the leading order spin-orbit interaction can be added to the PN accurate conservative Hamiltonian which gives the dynamics of nonspinning compact binaries.

It is highly desirable to obtain Keplerian-type parametric solution to the conservative dynamics of spinning compact binaries. This is so as 1PN accurate quasi-Keplerian parametrization, obtained by Damour and Deruelle [7], is heavily employed to analyze radio observations of relativistic binary pulsars and to test general relativity in strong field regimes [8, 9]. Moreover, 2PN accurate Keplerian-type parametric solution for nonspinning compact binaries, obtained in Refs. [10, 11], is heavily employed to construct highly efficient "ready to use" search templates for nonspinning compact binaries moving in inspiralling eccentric orbits [12].

Recently, 3PN accurate generalized quasi-Keplerian parametrization for the conservative dynamics of spinning compact binaries in eccentric orbits was obtained in Refs. [13, 14]. More precisely, the PN accurate parametric solutions are available for the following two distinct cases: (i) the binary is composed of two compact objects of equal mass and two arbitrary spins, and (ii) the two objects have unequal masses but only one object has a spin. The PN accurate parametric solution may be used to describe the following two astrophysical situations. As the measured neutron star (NS) masses in NS–NS binaries usually cluster around $1.4M_{\odot}$, case (i) may be used to describe such NS–NS binaries. The case (ii) may be applicable in the instance of a neutron star orbiting a much larger, rapidly rotating black hole. In that case, the black-hole spin will dominate the orbital precession, and the neutron star spin may be neglected.

The above parametrization, for the two cases considered, shows that the dynamics is integrable and cannot be chaotic [5]. The parametric solution will be employed to construct "ready to use" search templates of spinning compact binaries in inspiralling orbits [15]. Further, it will also be useful to construct a fully 2PN accurate "timing formula", required to analyze binary pulsar observations by the Square Kilometre Array (SKA) [16].

---

*Electronic address: `C.Koenigsdoerffer@uni-jena.de`
†Electronic address: `A.Gopakumar@uni-jena.de`

In this paper, we compute, using our parametrization, the observable dimensionless relativistic periastron advance parameter k, relevant for binary pulsars. The expression for k that we derive will be applicable, like the parametrization computed in Ref. [14], only for the two distinct cases mentioned above. Our expression for k is in agreement with the one obtained by Damour and Schäfer [10]. It should be noted that Damour and Schäfer employed the evolution equation for the Laplace-Runge-Lenz vector, computed from a Hamiltonian, to obtain the spin-orbit contributions to the parameter k. In this paper, we employ our generalized quasi-Keplerian parametrization, derived in Ref. [14], to compute the parameter k.

There are two reasons that motivated us to do the current computation. The 2PN accurate expression for k, derived by Damour and Schäfer, valid for arbitrary masses and spins, is extremely important due to the discovery of the double pulsar PSR J0737–3039 [17]. This is because the accurate timing of the double pulsar over a period of few years should lead to the measurement of the moment of inertia of one of the pulsars [18]. It is from the measurement of k to 2PN order, which includes leading order spin-orbit contributions, that one will be able to measure the moment of inertia of the fast rotating pulsar in this double pulsar [10]. The measurement of the moment of inertia of a neutron star for the first time will lead to strong constrains on its equation of state [19, 20]. We compute k, as mentioned earlier, using Keplerian-type parametrization with the aim of presenting an alternative way of obtaining this observable quantity.

A second reason for computing k is the existence of certain intriguing features appearing in our parametrization. We observed that the factor that modulates the true anomaly, due to spin-orbit interaction, contains a contribution, independent of the spins themselves, and of 1PN order [see Eq. (5.6k) in Ref. [14]]. We want to make sure that this feature will not appear in the observable parameter k, computed using our generalized quasi-Keplerian parametrization.

The paper is structured in the following manner. In Sec. II, we briefly describe the 3PN accurate Keplerian-type parametrization for the conservative dynamics of spinning compact binaries, moving in eccentric orbits, when spin effects are restricted to the leading order spin-orbit interaction, derived in Ref. [14], and point out its salient features. Sec. III and Sec. IV deal with the issues related to the derivation of the parameter k. We show for the first time how to derive k using the parametrization, described in Sec. II, after describing how Damour and Schäfer obtained k. We summarize our results in Sec. V. Appendices A and B provide details of our computation and consistency checks.

II. GENERALIZED QUASI-KEPLERIAN PARAMETRIZATION FOR SPINNING COMPACT BINARIES

Recently, 3PN accurate generalized quasi-Keplerian parametrization for compact binaries in eccentric orbits with the first-order spin-orbit interaction was obtained in Refs. [13, 14] for the following two distinct cases: (i) the binary is composed of two compact objects of equal mass and two arbitrary spins, and (ii) the two objects have unequal masses but only one object has a spin. The underlying dynamics is fully specified by the following PN accurate (reduced) Hamiltonian H, written symbolically as

$$\begin{aligned} H(\boldsymbol{r},\boldsymbol{p},\boldsymbol{S}_1,\boldsymbol{S}_2) &= H_{\rm N}(\boldsymbol{r},\boldsymbol{p}) + H_{\rm 1PN}(\boldsymbol{r},\boldsymbol{p}) + H_{\rm 2PN}(\boldsymbol{r},\boldsymbol{p}) \\ &\quad + H_{\rm 3PN}(\boldsymbol{r},\boldsymbol{p}) + H_{\rm SO}(\boldsymbol{r},\boldsymbol{p},\boldsymbol{S}_1,\boldsymbol{S}_2)\,, \end{aligned} \tag{2.1}$$

where $H_{\rm N}$, $H_{\rm 1PN}$, $H_{\rm 2PN}$, and $H_{\rm 3PN}$ are, respectively, the Newtonian, first, second and third PN contributions to the conservative dynamics of compact binaries, when the spin effects are neglected. The leading order spin-orbit coupling to the binary dynamics is given by $H_{\rm SO}$. In the above equation, $\boldsymbol{r} = \boldsymbol{\mathcal{R}}/(GM)$, $r = |\boldsymbol{r}|$, and $\boldsymbol{p} = \boldsymbol{\mathcal{P}}/\mu$, where $\boldsymbol{\mathcal{R}}$ and $\boldsymbol{\mathcal{P}}$ are the relative separation vector and its conjugate momentum vector, respectively. The familiar symbols M and μ have the usual meaning, namely the total mass and the reduced mass. The explicit expressions for PN corrections, associated with the motion of nonspinning compact binaries, were obtained in Refs. [10, 21]. The spin-orbit contribution, available in Refs. [10, 14, 22], reads

$$H_{\rm SO}(\boldsymbol{r},\boldsymbol{p},\boldsymbol{S}_1,\boldsymbol{S}_2) = \frac{1}{c^2 r^3}(\boldsymbol{r}\times\boldsymbol{p})\cdot\boldsymbol{S}_{\rm eff}\,, \tag{2.2}$$

where the effective spin $\boldsymbol{S}_{\rm eff}$ is defined by

$$\boldsymbol{S}_{\rm eff} \equiv \delta_1\boldsymbol{S}_1 + \delta_2\boldsymbol{S}_2\,, \tag{2.3}$$

with

$$\delta_1 = \frac{\eta}{2} + \frac{3}{4}\left(1 - \sqrt{1-4\eta}\right)\,, \tag{2.4a}$$

$$\delta_2 = \frac{\eta}{2} + \frac{3}{4}\left(1 + \sqrt{1-4\eta}\right)\,. \tag{2.4b}$$

The finite mass ratio η is given by $\eta = \mu/M$. The (reduced) spin vectors $\boldsymbol{S}_1$ and $\boldsymbol{S}_2$ are related to the individual spins $\boldsymbol{\mathcal{S}}_1$ and $\boldsymbol{\mathcal{S}}_2$ by $\boldsymbol{S}_1 = \boldsymbol{\mathcal{S}}_1/(\mu GM)$ and $\boldsymbol{S}_2 = \boldsymbol{\mathcal{S}}_2/(\mu GM)$, respectively. We recall that $\boldsymbol{\mathcal{R}}$, $\boldsymbol{\mathcal{P}}$, $\boldsymbol{\mathcal{S}}_1$, and $\boldsymbol{\mathcal{S}}_2$ are canonical variables, such that the orbital variables commute with the spin variables, e.g., see Refs. [6, 22].

In Ref. [14], we obtained the Keplerian-type parametrization in an orbital triad $(\boldsymbol{i},\boldsymbol{j},\boldsymbol{k})$. In this triad, the unit vector $\boldsymbol{i}$ is along the line of nodes, associated with the intersection of the orbital plane with the invariable plane $(\boldsymbol{e}_X,\boldsymbol{e}_Y)$, which is the plane perpendicular to the total (reduced) angular momentum $\boldsymbol{J}$, the only constant vector in our analysis. The unit vector $\boldsymbol{k}$ is defined by $\boldsymbol{k} \equiv \boldsymbol{L}/L$, where $\boldsymbol{L} = \boldsymbol{r} \times \boldsymbol{p}$ is the (reduced) orbital angular momentum vector, and hence $\boldsymbol{k}$ is perpendicular to the orbital plane (see Fig. 1). We display below, in its entirety, the parametric solution to the conservative 3PN accurate dynamics of spinning compact binaries moving in eccentric orbits, in ADM-type coordinates, with spin effects restricted to the leading order spin-orbit interactions. The dynamical vectors of the problem are given by

$$\boldsymbol{r}(t) = r(t)\cos\varphi(t)\,\boldsymbol{i}(t) + r(t)\sin\varphi(t)\,\boldsymbol{j}(t)\,, \tag{2.5}$$

$$\boldsymbol{L}(t) = L\boldsymbol{k}(t)\,, \tag{2.6}$$

$$\boldsymbol{S}(t) = J\boldsymbol{e}_Z - L\boldsymbol{k}(t)\,. \tag{2.7}$$

We note that in the equal-mass case (i) the (reduced) total spin $\boldsymbol{S}$ is the sum of two arbitrary spins $\boldsymbol{S}_1$ and $\boldsymbol{S}_2$, and in the single-spin case (ii) $\boldsymbol{S}$ is either $\boldsymbol{S}_1$ *or* $\boldsymbol{S}_2$. In our restricted cases, namely (i) and (ii), the (reduced) orbital angular momentum $\boldsymbol{L}$ and the (reduced) total spin $\boldsymbol{S}$ precess about the conserved (reduced) total angular momentum vector $\boldsymbol{J} = \boldsymbol{L} + \boldsymbol{S}$ at the same rate [14]. This configuration is called *simple precession* in the literature, e.g., see Ref. [23].

The time dependent basic vectors $(\boldsymbol{i}, \boldsymbol{j}, \boldsymbol{k})$ are explicitly given by

$$\boldsymbol{i}(t) = \cos\Upsilon(t)\boldsymbol{e}_X + \sin\Upsilon(t)\boldsymbol{e}_Y\,, \tag{2.8a}$$

$$\boldsymbol{j}(t) = -\cos\Theta\sin\Upsilon(t)\boldsymbol{e}_X + \cos\Theta\cos\Upsilon(t)\boldsymbol{e}_Y + \sin\Theta\,\boldsymbol{e}_Z\,, \tag{2.8b}$$

$$\boldsymbol{k}(t) = \sin\Theta\sin\Upsilon(t)\boldsymbol{e}_X - \sin\Theta\cos\Upsilon(t)\boldsymbol{e}_Y + \cos\Theta\,\boldsymbol{e}_Z\,. \tag{2.8c}$$

The angle Θ denotes the constant precession cone angle of $\boldsymbol{L} = L\boldsymbol{k}$ around $\boldsymbol{J}$ and is given by

$$\sin\Theta = \frac{S\sin\alpha}{J}\,, \tag{2.9a}$$

$$\cos\Theta = \frac{L + S\cos\alpha}{J}\,, \tag{2.9b}$$

where the magnitude of the total angular momentum is given by $J = (L^2 + S^2 + 2LS\cos\alpha)^{1/2}$ and α is the constant angle between $\boldsymbol{L}$ and $\boldsymbol{S}$.

The time evolution for $r(t)$, $\varphi(t)$, and $\Upsilon(t)$ is given in a parametric and PN accurate form, which reads

$$r = a_r\left(1 - e_r\cos u\right)\,, \tag{2.10a}$$

$$l \equiv n\left(t - t_0\right) = u - e_t\sin u + \left(\frac{g_{4t}}{c^4} + \frac{g_{6t}}{c^6}\right)(v - u) + \left(\frac{f_{4t}}{c^4} + \frac{f_{6t}}{c^6}\right)\sin v + \frac{i_{6t}}{c^6}\sin 2v + \frac{h_{6t}}{c^6}\sin 3v\,, \tag{2.10b}$$

$$\varphi - \varphi_0 = \left(1 + \tilde{k}\right)v + \left(\frac{f_{4\varphi}}{c^4} + \frac{f_{6\varphi}}{c^6}\right)\sin 2v + \left(\frac{g_{4\varphi}}{c^4} + \frac{g_{6\varphi}}{c^6}\right)\sin 3v + \frac{i_{6\varphi}}{c^6}\sin 4v + \frac{h_{6\varphi}}{c^6}\sin 5v\,, \tag{2.10c}$$

$$\Upsilon - \Upsilon_0 = \frac{\chi_{\rm so}J}{c^2L^3}(v + e\sin v)\,, \tag{2.10d}$$

$$\text{where}\quad v = 2\arctan\left[\left(\frac{1 + e_\varphi}{1 - e_\varphi}\right)^{1/2}\tan\frac{u}{2}\right]\,. \tag{2.10e}$$

The PN accurate expressions for the orbital elements a_r, e_r^2, n, e_t^2, $\tilde{k}$, and e_φ^2 and the PN orbital functions g_{4t}, g_{6t}, f_{4t}, f_{6t}, i_{6t}, h_{6t}, $f_{4\varphi}$, $f_{6\varphi}$, $g_{4\varphi}$, $g_{6\varphi}$, $i_{6\varphi}$, and $h_{6\varphi}$, in terms of E, L, S, η, and α, are available in Ref. [14]. The mass dependent coupling constant $\chi_{\rm so}$ connects the effective spin with the total (reduced) spin via $\boldsymbol{S}_{\rm eff} = \delta_1\boldsymbol{S}_1 + \delta_2\boldsymbol{S}_2 = \chi_{\rm so}\boldsymbol{S}$ and is given by

$$\chi_{\rm so} \equiv \begin{cases} \delta_1 = \delta_2 = 7/8 & \text{for (i), the equal-mass case}\,, \\ \delta_1(\eta) \text{ or } \delta_2(\eta) & \text{for (ii), the single-spin case}\,. \end{cases}$$

We display below the 3PN accurate expression for the *internal* parameter $\tilde{k}$, appearing in Eq. (2.10c) [this parameter was denoted by k in Ref. [14], see Eqs. (5.5c) and (5.6k) there]. It consists of two parts: the first one due to the 3PN accurate conservative dynamics of nonspinning point masses and the second one due to the leading order spin-orbit interaction

$$\tilde{k} = \tilde{k}_{\rm 3PN} + \tilde{k}_{\rm so}\,, \tag{2.11}$$

where

$$\tilde{k}_{\rm 3PN} = \frac{3}{c^2L^2}\left\{1 + \frac{(-2E)}{4c^2}\left(-5 + 2\eta + \frac{35 - 10\eta}{-2EL^2}\right) + \frac{(-2E)^2}{384c^4}\left[120 - 120\eta + 96\eta^2\right.\right.$$

$$+\frac{1}{(-2EL^2)}(-10080+13952\eta-123\pi^2\eta-1440\eta^2)$$

$$+\frac{1}{(-2EL^2)^2}(36960-40000\eta+615\pi^2\eta+1680\eta^2)\Bigg]\Bigg\}\,, \tag{2.12a}$$

$$\tilde{k}_{\rm so}=\frac{3}{c^2L^2}\left\{-\frac{\chi_{\rm so}}{3}-\chi_{\rm so}\frac{S}{L}\cos\alpha\right\}. \tag{2.12b}$$

It should be noted that $\tilde{k}_{\rm 3PN}$ gives the observable advance of periastron in the case of nonspinning compact binaries. This results from the construction of $\tilde{k}$ [see Sec. V in Ref. [14] for details and Ref. [10]]. However, when spin-orbit interactions are added to the dynamics of compact binaries, $\tilde{k}$ does not directly lead to the measurable advance of periastron, relevant for binary pulsars. This parameter $\tilde{k}$ is a measure for the advance of periastron *only inside* our parametrization and has to be related to the observational one. Further, note the presence of the $\chi_{\rm so}$ term (which does not depend on the spins), appearing in $\tilde{k}_{\rm so}$ at the 1PN order. It is due to these points, namely the $\chi_{\rm so}$ term that appears at the 1PN order and $\tilde{k}$ does not give the observable periastron advance for spinning compact binaries, that we compute the dimensionless observable periastron advance parameter k for spinning compact binaries in this paper. Note, that we have corrected the sign in front of $\chi_{\rm so}/3$ in $\tilde{k}_{\rm so}$, important for the proper nonspinning limit and the following calculation of the observable periastron advance parameter k in Sec. IV.

Before we go onto the details of computing k, let us list a few salient featurs of the parametrization. The parameter e, appearing in Eq. (2.10d), is given by the Newtonian contribution to say e_r. Note that at Newtonian order all three eccentricities e_r, e_t, and e_φ, are identical to each other and at PN orders they are related by PN accurate relations [see Eqs. (5.7) in Ref. [14]]. Therefore, it is possible to describe the binary dynamics in terms of one of the eccentricities. We emphasize that, while adapting the parametrization to describe the binary dynamics, care should be taken to restrict orbital elements to the required PN order. We are aware that the above parametrization does not lead to the classic Keplerian parametrization simply by putting $S=0$. The detailed explanation of the nonspinning limit and the apparent lack of a *simple* nonspinning limit is given in Sec. IV B in Ref. [14].

The above parametric solution is usually refered to as the genaralized quasi-Keplerian parametrization for the PN accurate orbital dynamics of spinning compact binaries. It extends the computations of Damour, Deruelle, Schäfer, and Wex [7, 10, 11, 13, 24].

Finally, we want to emphasize that the parametric solution, given by Eqs. (2.5)–(2.12), describes the entire conservative PN accurate dynamics of a spinning compact binary in an eccentric orbit, when the spin effects are restricted to the leading order spin-orbit interaction. This means the parametrization consistently describes not only the precessional motion of the orbit inside the orbital plane, but also the precessional motions of the orbital plane and the spins themselves. Further, the deterministic nature of the dynamics described by the Hamiltonian, given by Eq. (2.1) and treated as a self-contained dynamical system, naturally follows from the above parametric solution.

In the next section, we develop the basics required to compute k from our parametrization.

III. RELEVANT INPUTS REQUIRED TO COMPUTE k

In this section, we will introduce two new triads, in terms of which it will be easy to obtain k. We will also discuss in detail how to connect these triads and hence the parametrization expressed in terms of these triads.

A. The periastron-based triad for the parametrization

As we are interested in computing k using our parametrization, it is natural that we introduce a periastron-based triad $(\boldsymbol{a},\boldsymbol{b},\boldsymbol{k})$. Later, we will express the relative separation vector $\boldsymbol{r}$, given by Eq. (2.5), in terms of this new triad.

Let us introduce this periastron-based triad in a pedagogical way. We start by writing the angle φ, given by Eq. (2.10c), in the following way

$$\varphi=v+\Pi\,. \tag{3.1}$$

Remember, φ is the polar angle inside the orbital plane measured from the line of nodes $\boldsymbol{i}$, associated with the intersection of the orbital plane with the invariable plane $(\boldsymbol{e}_X,\boldsymbol{e}_Y)$, and v is the "true anomaly", i.e., the angle between the (instantaneous) periastron and the position of the relative separation vector $\boldsymbol{r}$. The angle Π may be considered as the "argument of the periastron", i.e., the angle between the line of nodes $\boldsymbol{i}$ and the (instantaneous) periastron. By comparing Eqs. (3.1) and (2.10c), we can easily infer the parametric evolution for Π. Therefore, it is obvious that the angle Π contains via $\tilde{k}$ the secular contributions to the periastron advance.

Inserting Eq. (3.1) into Eq. (2.5) and making use of



Figure 1: The binary geometry and definition of various angles. Our reference frame is $(\boldsymbol{e}_X, \boldsymbol{e}_Y, \boldsymbol{e}_Z)$, where the basic vector $\boldsymbol{e}_Z$ is aligned with the fixed total angular momentum vector $\boldsymbol{J} = \boldsymbol{L} + \boldsymbol{S}$, such that $\boldsymbol{J} = J\boldsymbol{e}_Z$. The invariable plane $(\boldsymbol{e}_X,\ \boldsymbol{e}_Y)$ is perpendicular to $\boldsymbol{J}$. Important for the observation is the line-of-sight unit vector $\boldsymbol{N}$, pointing from the source (compact binary) to the observer. We may have, by a clever choice of $\boldsymbol{e}_X$ and $\boldsymbol{e}_Y$, the line-of-sight unit vector $\boldsymbol{N}$ in the $\boldsymbol{e}_Y$-$\boldsymbol{e}_Z$ plane. $\boldsymbol{k} = \boldsymbol{L}/L$ is the unit vector in the direction of the orbital angular momentum $\boldsymbol{L}$, which is perpendicular to the orbital plane. The constant inclination of the orbital plane with respect to the invariable plane is Θ, which is also the precession cone angle of $\boldsymbol{L}$ around $\boldsymbol{J}$. The three lines of nodes are associated with the intersections of the orbital plane, the invariable plane and the plane of the sky. The orbital plane intersects the invariable plane at the line of nodes $\boldsymbol{i}$, with longitude Υ, measured in the invariable plane from $\boldsymbol{e}_X = \boldsymbol{p}$, which also gives the phase of the orbital plane precession. The orbital plane intersects the plane of the sky at the line of nodes $\boldsymbol{i}_i$, with longitude Ω, measured in the plane of the sky, from $\boldsymbol{e}_X = \boldsymbol{p}$. The subscript i in $\boldsymbol{i}_i$ results from the corresponding *orbital* inclination angle i. The *dynamical* longitude of periastron, measured in the orbital plane from $\boldsymbol{i}$, is Π. The *observational* longitude of periastron, measured in the orbital plane from $\boldsymbol{i}_i$, is ω. The above figure implies $\omega = \Delta + \Pi$.

familiar trigonometric relations leads to

$$\begin{aligned} \boldsymbol{r} &= r\cos(v+\Pi)\,\boldsymbol{i} + r\sin(v+\Pi)\,\boldsymbol{j} \\ &= r\cos v\,\boldsymbol{a} + r\sin v\,\boldsymbol{b}\,, \end{aligned} \tag{3.2}$$

where the new basic vectors are given by

$$\boldsymbol{a} = \cos\Pi\,\boldsymbol{i} + \sin\Pi\,\boldsymbol{j}\,, \tag{3.3a}$$

$$\boldsymbol{b} = -\sin\Pi\,\boldsymbol{i} + \cos\Pi\,\boldsymbol{j}\,. \tag{3.3b}$$

Now, it is obvious that $\boldsymbol{a}$ is the unit vector pointing towards the (instantaneous) periastron, as $\boldsymbol{r}$ is periodic in the angular variable v with period 2π in these new basic vectors $\boldsymbol{a}$ and $\boldsymbol{b}$. Hence it is straightforward to define the periastron-based orthonormal triad $(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{k}) \equiv (\boldsymbol{a}, \boldsymbol{k}\times\boldsymbol{a}, \boldsymbol{k})$, by rotating the $(\boldsymbol{i}, \boldsymbol{j}, \boldsymbol{k})$ frame around the orbital angular momentum unit vector $\boldsymbol{k}$, by an angle Π.

This gives us

$$\begin{pmatrix} \boldsymbol{a} \\ \boldsymbol{b} \\ \boldsymbol{k} \end{pmatrix} = \begin{pmatrix} \cos\Pi & \sin\Pi & 0 \\ -\sin\Pi & \cos\Pi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \boldsymbol{i} \\ \boldsymbol{j} \\ \boldsymbol{k} \end{pmatrix}\,, \tag{3.4}$$

where Π is the angle between $\boldsymbol{i}$ and the (instantaneous) periastron unit vector $\boldsymbol{a}$, as explained above.

The periastron-based vectors $(\boldsymbol{a}, \boldsymbol{b})$ are co-rotating with the precessing (instantaneous) Newtonian ellipse. This is similar to the behaviour of a body fixed reference system usually encountered in classical mechanics [25]. Hence, the set of angles (Υ, Π, Θ) is equivalent to the Eulerian angles defined there.

In the next subsection, we will introduce an observer-based triad and investigate how to connect the observer and periastron-based triads.

B. Connecting the observer and periastron-based triads

The second new triad that we introduce is a fixed one associated with the observer on earth and denoted by $(\boldsymbol{p},\boldsymbol{q},\boldsymbol{N})$. In this fixed triad $\boldsymbol{p}$ and $\boldsymbol{q}$ are two orthogonal unit vectors in the plane of the sky, i.e., the plane transverse to the radial direction $\boldsymbol{N}$, pointing from the source (compact binary) to the observer (see Fig. 1).

In order to compute k using our parametrization, it is natural to expect that we need to connect the periastron-based triad $(\boldsymbol{a},\boldsymbol{b},\boldsymbol{k})$ to the observer triad $(\boldsymbol{p},\boldsymbol{q},\boldsymbol{N})$. It is possible to connect these two frames with the help of rotational matrices quite similar to the way in which the $(\boldsymbol{i},\boldsymbol{j},\boldsymbol{k})$ frame was connected to the $(\boldsymbol{p},\boldsymbol{q},\boldsymbol{N})$ frame in Ref. [14] [see Sec. VI in Ref. [14] for details]. In this paper, we come from the observer triad $(\boldsymbol{p},\boldsymbol{q},\boldsymbol{N})$ to the periastron-based triad $(\boldsymbol{a},\boldsymbol{b},\boldsymbol{k})$ with the help of *four* successive rotations: first, a rotation around $\boldsymbol{p}$ via the *constant* angle i_0 to come to the reference triad $(\boldsymbol{e}_X,\boldsymbol{e}_Y,\boldsymbol{e}_Z)$, second, a rotation around $\boldsymbol{e}_Z$ by an angle Υ, third, a rotation around the direction of the line of nodes $\boldsymbol{i}$ [associated with the intersection of the orbital plane with the invariable plane $(\boldsymbol{e}_X,\boldsymbol{e}_Y)$] by an angle Θ, and, finally fourth, a rotation around the orbital angular momentum direction $\boldsymbol{k}$ by an angle Π. In terms of rotational matrices — denoted by $\mathcal{R}$[rotation angle, rotation axis] — we have

$$\begin{pmatrix}\boldsymbol{a}(t)\\ \boldsymbol{b}(t)\\ \boldsymbol{k}(t)\end{pmatrix} = \mathcal{R}[\Pi(t),\boldsymbol{k}(t)]\;\mathcal{R}[\Theta,\boldsymbol{i}(t)]\;\mathcal{R}[\Upsilon(t),\boldsymbol{e}_Z] \times \mathcal{R}[i_0,\boldsymbol{p}]\begin{pmatrix}\boldsymbol{p}\\ \boldsymbol{q}\\ \boldsymbol{N}\end{pmatrix}, \qquad (3.5)$$

where the first rotation appears at the right-hand side, the second one appears left from the one before, and so on.

We note, that the approach to generate the $(\boldsymbol{a},\boldsymbol{b},\boldsymbol{k})$ frame via these rotations may be also adapted for scenarios, where the angle Θ depends on time. The constant behaviour of Θ is a direct consequence of choosing two distinct cases (i) and (ii), where our parametrization is valid and hence we will use the fact that Θ is constant only at the end of our analysis.

It should be noted that there are other possible ways to connect the orthonormal triads $(\boldsymbol{p},\boldsymbol{q},\boldsymbol{N})$ and $(\boldsymbol{a},\boldsymbol{b},\boldsymbol{k})$. A second way to generate the $(\boldsymbol{a},\boldsymbol{b},\boldsymbol{k})$ frame is described in great detail by Damour and Schäfer in Sec. 5 of Ref. [10] and involves *three* rotational matrices. This way is closely linked to the observation as it involves rotational matrices defined with the help of the observational longitude of periastron ω and the orbital inclination angle i. Damour and Schäfer connected the observer triad $(\boldsymbol{p},\boldsymbol{q},\boldsymbol{N})$ to the periastron-based triad $(\boldsymbol{a},\boldsymbol{b},\boldsymbol{k})$ via the following *three* successive rotations: first, a rotation around $\boldsymbol{N}$ by an angle Ω, second, a rotation around the direction of the line of nodes $\boldsymbol{i}_i$ [associated with the intersection of the orbital plane with the plane of the sky $(\boldsymbol{p},\boldsymbol{q})$] by an angle i, and, finally third, a rotation around the orbital angular momentum direction $\boldsymbol{k}$ by an angle ω. The corresponding rotational matrices read

$$\begin{pmatrix}\boldsymbol{a}(t)\\ \boldsymbol{b}(t)\\ \boldsymbol{k}(t)\end{pmatrix} = \mathcal{R}[\omega(t),\boldsymbol{k}(t)]\;\mathcal{R}[i(t),\boldsymbol{i}_i(t)]\;\mathcal{R}[\Omega(t),\boldsymbol{N}]\begin{pmatrix}\boldsymbol{p}\\ \boldsymbol{q}\\ \boldsymbol{N}\end{pmatrix}. \qquad (3.6)$$

We note, that for the unit vector $\boldsymbol{i}_i$ along the line of nodes, associated with the intersection of the orbital plane with the plane of the sky, the subscript i is employed as $\boldsymbol{i}_i$ is related to the orbital inclination angle i and to avoid any confusion with the unit vector $\boldsymbol{i}$.

In the next section, we obtain k with the help of the parametrization and the ideas discussed in this section.

IV. THE ADVANCE OF PERIASTRON RELEVANT FOR BINARY PULSARS

In this section, we will compute k, the observable dimensionless periastron advance parameter due to $\tilde{k}$. Following Damour and Schäfer, the PN accurate total observable dimensionless periastron advance parameter k can be written as

$$k = k_{\rm 3PN} + k_{\rm so}\,, \qquad (4.1)$$

where $k_{\rm 3PN}$ is the usual PN accurate periastron advance parameter up to 3PN order, corresponding to the non-spinning case [see Refs. [10, 13] for the 2PN and 3PN contributions, respectively], and $k_{\rm so}$ denotes the contribution due to spin-orbit interaction. It should be kept in mind that the 3PN contribution to $k_{\rm 3PN}$ will not be required while probing the dynamics of relativistic binary pulsars even with SKA.

Let us first briefly summarize how Damour and Schäfer obtained k, especially $k_{\rm so}$, in the next subsection.

A. The derivation of $k_{\rm so}$ — The approach of Damour and Schäfer

Timing of binary pulsars allows via the observed “Roemer time delay” the direct measurement of the argument of periastron ω [8]. Therefore, Damour and Schäfer defined the relevant *secular* periastron advance as $\langle d\omega/dt\rangle_t$, or equivalently the timing parameter (not to be confused with the orbital angular-momentum unit vector $\boldsymbol{k}$)

$$k \equiv \frac{1}{n}\left\langle\frac{d\omega}{dt}\right\rangle_t\,, \qquad (4.2)$$

where n is the mean orbital motion that appears in the PN accurate Kepler equation, given by Eq. (2.10b), and $\langle\ldots\rangle_t$ denotes the orbital average.

Damour and Schäfer computed the contribution to $d\omega/dt$ due to spin-orbit interaction in the following way.

First, they computed the precessional equations for the orbital angular momentum vector $\boldsymbol{\mathcal{L}}$ and the Laplace-Runge-Lenz vector $\boldsymbol{\mathcal{A}}$, which tracks the periastron advance, with the help of the 2PN accurate version of Eq. (2.1). These precessional equations where orbital averaged to obtain the rotational velocity vector, denoted by $\boldsymbol{\Omega}$ in Ref. [10]. The expression for $d\omega/dt$ was extracted from $\boldsymbol{\Omega}$ by decomposing $\boldsymbol{\Omega}$ along three *non-orthogonal* directions [see Eq. (5.11) in Ref. [10]]. In this way a parametric solution to the dynamics of spinning compact binaries was not required to compute the 2PN accurate measurable dimensionless periastron advance parameter k in Ref. [10], and hence the formula of Damour and Schäfer for k is more general.

In the next subsection, we will compute $d\omega/dt$ using our Keplerian-type parametric solution and the two different ways of connecting the $(\boldsymbol{p}, \boldsymbol{q}, \boldsymbol{N})$ frame to the $(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{k})$ frame.

B. The derivation of k_{so} — The parametric approach

This subsection deals with the computation of $d\omega/dt$, arising due to spin-orbit interaction, using our Keplerian-type parametrization.

Following Damour and Schäfer, we introduce the rotation velocity vector $\boldsymbol{\Omega}$, containing all involved angular velocities, and extract $d\omega/dt$ from it in the following way. We have already noted that there are two different ways to connect the observer-based triad to the pariastron-based one [see Eqs. (3.5) and (3.6) and related discussions there]. The first way, given by Eq. (3.5), involves the angular variables $\Pi(t)$, Θ, $\Upsilon(t)$, and i_0, appearing in our parametrization, and the second way, given by Eq. (3.6), involves the angles $\omega(t)$, $i(t)$, and $\Omega(t)$, where two of them are relevant for the observations, namely $\omega(t)$ and $i(t)$. Therefore, we can decompose the rotation velocity vector $\boldsymbol{\Omega}$ in the following two ways:

$$\boldsymbol{\Omega} = \frac{d\Pi}{dt}\boldsymbol{k} + \frac{d\Theta}{dt}\boldsymbol{i} + \frac{d\Upsilon}{dt}\boldsymbol{e}_Z + \frac{di_0}{dt}\boldsymbol{p}\,, \tag{4.3a}$$

$$\boldsymbol{\Omega} = \frac{d\omega}{dt}\boldsymbol{k} + \frac{di}{dt}\boldsymbol{i}_i + \frac{d\Omega}{dt}\boldsymbol{N}\,. \tag{4.3b}$$

Note, that the above two different decompositions are along four/three *non-orthogonal* directions, respectively. The elementary rotation velocities, appearing in Eq. (4.3a), can be defined with the help of our parametrization, whereas in Eq. (4.3b) we have $d\omega/dt$: the rate of change of the argument of the periastron. Further note, that Eq. (4.3b) is identical to Eq. (5.11) of Ref. [10].

In order to extract $d\omega/dt$ and to relate it to the angular variables associated with our parametrization we proceed as follows. Since all vectors, appearing in Eqs. (4.3), can be expressed in terms of $(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{k})$, it is clear that the rotation velocity vector $\boldsymbol{\Omega}$ may also be written as the sum of three elementary rotation velocities, along three *orthogonal* directions $\boldsymbol{a}$, $\boldsymbol{b}$, and $\boldsymbol{k}$, namely

$$\boldsymbol{\Omega} = \Omega_{(\boldsymbol{a})}\boldsymbol{a} + \Omega_{(\boldsymbol{b})}\boldsymbol{b} + \Omega_{(\boldsymbol{k})}\boldsymbol{k}\,. \tag{4.4}$$

It is then straightforward to compute the components $\Omega_{(\boldsymbol{a})}$, $\Omega_{(\boldsymbol{b})}$, and $\Omega_{(\boldsymbol{k})}$ in terms of the elementary rotation velocities and angles appearing in Eqs. (4.3). The components of $\boldsymbol{\Omega}$ in the co-rotating periastron-based $(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{k})$ frame — using the parametrization-based variables and the observation-based variables, respectively — are given by

$$\Omega_{(\boldsymbol{a})} = \frac{d\Theta}{dt}\cos\Pi + \frac{d\Upsilon}{dt}\sin\Theta\sin\Pi + \frac{di_0}{dt}(\cos\Pi\cos\Upsilon - \cos\Theta\sin\Pi\sin\Upsilon) = \frac{di}{dt}\cos\omega + \frac{d\Omega}{dt}\sin i\sin\omega\,, \tag{4.5a}$$

$$\Omega_{(\boldsymbol{b})} = -\frac{d\Theta}{dt}\sin\Pi + \frac{d\Upsilon}{dt}\sin\Theta\cos\Pi + \frac{di_0}{dt}(-\sin\Pi\cos\Upsilon - \cos\Theta\cos\Pi\sin\Upsilon) = -\frac{di}{dt}\sin\omega + \frac{d\Omega}{dt}\sin i\cos\omega\,, \tag{4.5b}$$

$$\Omega_{(\boldsymbol{k})} = \frac{d\Pi}{dt} + \frac{d\Upsilon}{dt}\cos\Theta + \frac{di_0}{dt}\sin\Theta\sin\Upsilon = \frac{d\omega}{dt} + \frac{d\Omega}{dt}\cos i\,. \tag{4.5c}$$

These *general* set of equations provides the transformation from the parametrized angles to the angles of interest for the observation. We note, that the last component, $\Omega_{(\boldsymbol{k})}$, gives the so-called *intrinsic* contribution to the periastron advance $\dot{\omega}^{\text{intrinsic}}$, defined by $\dot{\omega}^{\text{intrinsic}} \equiv \boldsymbol{\Omega}\cdot\boldsymbol{k}$. This is the component of $\boldsymbol{\Omega}$ acting *within* the orbital plane. However, in the timing of binary pulsars the parameter k enters in the (relativistic) argument of the periastron ω, and so we have to extract $d\omega/dt$ from the above set of equations.

A close inspection of Fig. 1 indicates that i_0 is constant, since it connects two invariant reference frames. For the two distinct cases (i) and (ii), where our parametrization is valid, we also have Θ as a constant angle. This allows us to come from Eqs. (4.5) to the *reduced* set of equations for the components $\Omega_{(\boldsymbol{a})}$, $\Omega_{(\boldsymbol{b})}$, and $\Omega_{(\boldsymbol{k})}$, which reads

$$\frac{d\Upsilon}{dt}\sin\Theta\sin\Pi = \frac{di}{dt}\cos\omega + \frac{d\Omega}{dt}\sin i\sin\omega\,, \tag{4.6a}$$

$$\frac{d\Upsilon}{dt}\sin\Theta\cos\Pi = -\frac{di}{dt}\sin\omega + \frac{d\Omega}{dt}\sin i\cos\omega\,, \tag{4.6b}$$

$$\frac{d\Pi}{dt}+\frac{d\Upsilon}{dt}\cos\Theta=\frac{d\omega}{dt}+\frac{d\Omega}{dt}\cos i\,. \tag{4.6c}$$

Solving the last equation, namely Eq. (4.6c), for $d\omega/dt$ leads to

$$\frac{d\omega}{dt}=\frac{d\Pi}{dt}+\frac{d\Upsilon}{dt}\cos\Theta-\frac{d\Omega}{dt}\cos i\,. \tag{4.7}$$

The first and the second contribution to $d\omega/dt$ are directly provided by our parametrization. We also figured out that they can be obtained via the time evolution of the Laplace-Runge-Lenz vector without any parametrization (see Appendix B for details). The third contribution to $d\omega/dt$ in Eq. (4.7), namely $(d\Omega/dt)\cos i$, needs a closer inspection. To obtain $d\Omega/dt$, we multiply Eq. (4.6a) with $\sin\omega$ and add it to the product of Eq. (4.6b) times $\cos\omega$, which results in

$$\frac{d\Omega}{dt}=\frac{1}{\sin i}\frac{d\Upsilon}{dt}\sin\Theta\left(\sin\Pi\sin\omega+\cos\Pi\cos\omega\right)\,. \tag{4.8}$$

Using now the explicit expressions for $\sin\omega$ and $\cos\omega$, given by Eqs. (A9) in Appendix A, in the above equation, we get

$$\frac{d\Omega}{dt}=\frac{1}{\sin^2 i}\frac{d\Upsilon}{dt}S_\Theta(C_{i_0}S_\Theta+S_{i_0}C_\Theta\cos\Upsilon)\,, \tag{4.9}$$

where S_Θ, C_Θ, S_{i_0}, and C_{i_0} are shorthand notations for the constant trigonometric functions $\sin\Theta$, $\cos\Theta$, $\sin i_0$, and $\cos i_0$, respectively. The time evolution for the orbital inclination i, appearing in Eqs. (4.7)–(4.9), is simply given by Eqs. (A4) in Appendix A.

Let us now explicitly compute $d\Pi/dt$, $d\Upsilon/dt$ and hence $d\Omega/dt$ with the help of our parametric solution. Using Eqs. (2.10c), (2.11), and (3.1) for Π, and Eq. (2.10d) for Υ, we display below the relevant parametric expressions symbolically

$$\begin{aligned}\Pi-\Pi_0&=\left(\tilde{k}_{\rm 3PN}+\tilde{k}_{\rm so}\right)v+(\ldots)\sin 2v+(\ldots)\sin 3v\\&\quad+(\ldots)\sin 4v+(\ldots)\sin 5v\,, &(4.10a)\\ \Upsilon-\Upsilon_0&=\frac{\chi_{\rm so}J}{c^2L^3}(v+e\sin v)\,. &(4.10b)\end{aligned}$$

The structure of Eqs. (4.10) indicates that we can handle the PN and the spin-orbit induced contributions to $d\omega/dt$ separately.

To obtain the spin-orbit contributions to $\langle d\omega/dt\rangle_t$, we compute with the help of Eqs. (4.10) and (2.12b) the orbital averaged spin-orbit (so) contributions to the above required time derivatives

$$\begin{aligned}\left\langle\frac{d\Pi}{dt}\right\rangle_t^{\rm so}&=\frac{1}{c^2}\left\langle\frac{1}{r^3}\right\rangle_t\chi_{\rm so}(-L-3S\cos\alpha)\,, &(4.11a)\\ \left\langle\frac{d\Upsilon}{dt}C_\Theta\right\rangle_t^{\rm so}&=\frac{1}{c^2}\left\langle\frac{1}{r^3}\right\rangle_t\chi_{\rm so}(L+S\cos\alpha)\,, &(4.11b)\\ \left\langle\frac{d\Omega}{dt}\cos i\right\rangle_t^{\rm so}&=\frac{1}{c^2}\left\langle\frac{1}{r^3}\right\rangle_t\chi_{\rm so}S\sin\alpha\frac{\cos i}{\sin^2 i}\\&\quad\times(C_{i_0}S_\Theta+S_{i_0}C_\Theta\cos\Upsilon)\,. &(4.11c)\end{aligned}$$

The quantity $\langle 1/r^3\rangle_t$ gives the orbital average of $1/r^3$, r being the scaled radial separation, to the Newtonian order [see, e.g., Ref. [10]]

$$\left\langle\frac{1}{r^3}\right\rangle_t=\frac{1}{a_r^3(1-e^2)^{3/2}}=\frac{n^2}{(1-e^2)^{3/2}}\,. \tag{4.12}$$

With the help of Eqs. (4.7), (4.11), and (4.12), it is straightforward to obtain the spin-orbit induced contribution to k, defined by Eqs. (4.1) and (4.2), and it reads

$$k_{\rm so}=B\left[-2\cos\alpha-\sin\alpha\frac{\cos i}{\sin^2 i}(C_{i_0}S_\Theta+S_{i_0}C_\Theta\cos\Upsilon)\right]\,, \tag{4.13}$$

where B is given in terms of our *reduced* variables by

$$B=\frac{n\chi_{\rm so}S}{c^2(1-e^2)^{3/2}}\,. \tag{4.14}$$

The proper nonspinning limit of Eqs. (4.13) and (4.14), can be easily deduced as when $S\to 0$, $\boldsymbol{J}$ is identical to $\boldsymbol{L}$, which implies $\Theta\to 0$ and $i_0\to i$, i being the angle between $\boldsymbol{N}$ and $\boldsymbol{L}$, and therefore $k_{\rm so}$ vanishes. The disappearance of $k_{\rm so}$ as $S\to 0$ indicates that the presence of the 1PN accurate $\chi_{\rm so}$ term in $\tilde{k}_{\rm so}$, given by Eq. (2.12b), does not have any observational consequence.

Our result for $k_{\rm so}$ is in agreement with Eqs. (5.17) and (5.17a) of Ref. [10], when we specialize these equations to the *simple precession* (for the equal-mass case and for the single-spin case)

$$k_{\rm so}=\frac{B}{\sin^2 i}\left[(1-3\sin^2 i)(\boldsymbol{k}\cdot\boldsymbol{s})-\cos i(\boldsymbol{N}\cdot\boldsymbol{s})\right]\,, \tag{4.15}$$

where $\boldsymbol{s}\equiv\boldsymbol{\mathcal{S}}/\mathcal{S}$, $\boldsymbol{k}\cdot\boldsymbol{s}=\cos\alpha$ and the coefficient B is given here in terms of the *non-reduced* variables, used by Damour and Schäfer in Ref. [10] and in Appendix B, by

$$B=\frac{n\lambda_{\rm so}\mathcal{S}}{c^2M(1-e^2)^{3/2}}\,. \tag{4.16}$$

The dimensionless mass dependent coupling constant $\lambda_{\rm so}$ is related to $\chi_{\rm so}$ by $\lambda_{\rm so}=\chi_{\rm so}/\eta$ (see Appendix B for details). To check the above agreement between our result and the one of Damour and Schäfer, we have used Eqs. (2.9) and Eq. (A4a). The two expressions for the coefficient B, namely Eqs. (4.14) and (4.16), are identical.

Let us now briefly consider the nonspinning 3PN accurate contributions to k. These contributions, enter only via Π, orbital averaged analogue to Eq. (4.11a), our computation, and result in $k_{\rm 3PN}\equiv\tilde{k}_{\rm 3PN}$. However, it should be noted that the expression for $k_{\rm 3PN}$, as given by Eq. (2.12a), is not yet ready for application. This is so as this expression for $k_{\rm 3PN}$ depends on the conserved *theoretical* quantities E and L rather than on *observable* quantities like mean motion n and proper-time eccentricity e_T. However, we have checked that up to 2PN order our expression for $k_{\rm 3PN}$ is in agreement with the one of Damour and Schäfer, obtained in Ref. [10].

Therefore, the 3PN accurate observable periastron advance parameter k becomes

$$k = k_{\rm 3PN} + k_{\rm so}\,, \tag{4.17}$$

where $k_{\rm 3PN}$ is the *orbital* 3PN contribution, corresponding to the nonspinning case, given by the 3PN accurate contribution to Eq. (2.11), namely Eq. (2.12a). The spin-orbit contribution $k_{\rm so}$ is given by Eqs. (4.13) and (4.14) or equivalently by Eqs. (4.15) and (4.16). We recapitulate that $k_{\rm 3PN} \equiv \tilde{k}_{\rm 3PN}$ and $k_{\rm so} \neq \tilde{k}_{\rm so}$, where $\tilde{k}_{\rm 3PN}$ and $\tilde{k}_{\rm so}$ are the corresponding contributions to $\tilde{k}$, as given by Eqs. (2.11) and (2.12). This results from the construction of $\tilde{k}$ in Ref. [14] by combining the parameter $\tilde{k}_{\rm 3PN} = k_{\rm 3PN}$ of the nonspinning case with the internal parameter $\tilde{k}_{\rm so}$, given by Eq. (4.45e) in Ref. [14], of the spin-orbit parametrization [see Sec. V in Ref. [14] for details].

Finally, let us present the following geometric explanation why $\tilde{k} \neq k$. The difference between $\tilde{k}$ and k results from the fact, that the angle Π — associated with the parameter $\tilde{k}$ — is measured from $\boldsymbol{i}$ and in contrast to that, the angle ω — associated with the observable parameter k — is measured with respect to $\boldsymbol{i}_i$. The difference between ω and Π is denoted by Δ in Fig. 1 (see Appendix A for details). In addition, we want to point out, that the last rotation $\mathcal{R}[\omega(t), \boldsymbol{k}(t)]$ in Eq. (3.6) can be decomposed into two parts

$$\mathcal{R}[\omega(t), \boldsymbol{k}(t)] = \mathcal{R}[\Pi(t), \boldsymbol{k}(t)]\; \mathcal{R}[\Delta(t), \boldsymbol{k}(t)]\,, \tag{4.18}$$

and so it is obvious that $\omega(t) = \Delta(t) + \Pi(t)$ and correspondingly we have $d\omega/dt = d\Delta/dt + d\Pi/dt$. The contribution of $d\Delta/dt$ causes the difference between $k_{\rm so}$ and $\tilde{k}_{\rm so}$.

V. CONCLUSIONS

We have presented a new method to compute the PN accurate observable dimensionless relativistic periastron advance parameter k of a spinning compact binary using our Keplerian-type parametric solution to the PN accurate conservative dynamics of spinning compact binaries moving in eccentric orbits [14]. The expression for k is applicable, like our parametric solution, only for the following two distinct cases, namely, case (i), the equal-mass case and case (ii), the single-spin case. Our result is consistent with a more general formula for k, derived by Damour and Schäfer, using a different procedure [10]. The computation also helped us to clarify certain intriguing features that we observed in our parametrization.

Acknowledgments

It is our pleasure to thank Gerhard Schäfer for helpful discussions and encouragements. We gratefully acknowledge the financial support of the Deutsche Forschungsgemeinschaft (DFG) through SFB/TR7 “Gravitationswellenastronomie”.

The algebraic computations, appearing in this paper, were performed using Maple and Mathematica.

Appendix A: SOME ASPECTS OF THE BINARY GEOMETRY AND RELATED DYNAMICS

In this section, we make contact with the variables and angles, commonly used in the literature [10, 26]. It is obvious that we have three planes in the binary geometry, namely the orbital plane, the invariable plane and the plane of the sky. The associated orthonormal triads are $(\boldsymbol{i}, \boldsymbol{j}, \boldsymbol{k})$, $(\boldsymbol{e}_X, \boldsymbol{e}_Y, \boldsymbol{e}_Z)$, and $(\boldsymbol{p}, \boldsymbol{q}, \boldsymbol{N})$. Therefore, we may define three lines of nodes and associated inclination angles in the following way. We infer from Fig. 1 that the plane of the sky meets the invariable plane along a line of nodes, characterized by $\boldsymbol{p} = \boldsymbol{e}_X$, as we choose the line-of-sight unit vector $\boldsymbol{N}$ to lie in the $\boldsymbol{e}_Y$-$\boldsymbol{e}_Z$ plane. The associated constant angle i_0 between $\boldsymbol{N}$ and $\boldsymbol{J} = J\boldsymbol{e}_Z$ may be treated as the inclination angle, related to the above line of nodes. In the second case, we consider the line of nodes $\boldsymbol{i}$, created by the intersection of the orbital plane with the invariable plane. The corresponding inclination angle, namely the angle between the orbital angular momentum unit vector $\boldsymbol{k} = \boldsymbol{L}/L$ and $\boldsymbol{J} = J\boldsymbol{e}_Z$, is denoted by Θ. Finally, we consider the intersection of the orbital plane with the plane of the sky. We associate $\boldsymbol{i}_i$ with this line of nodes and the related inclination angle i, the angle between $\boldsymbol{k}$ and the line of sight unit vector $\boldsymbol{N}$, is the usually defined orbital inclination i and hence the use of the subscript i in $\boldsymbol{i}_i$.

It is easy to figure out that the unit vectors, associated with the last two lines of nodes, $\boldsymbol{i}$ and $\boldsymbol{i}_i$ are defined by

$$\boldsymbol{i} \equiv \frac{\boldsymbol{e}_Z \times \boldsymbol{k}}{|\boldsymbol{e}_Z \times \boldsymbol{k}|}\,, \tag{A1a}$$

$$\boldsymbol{i}_i \equiv \frac{\boldsymbol{N} \times \boldsymbol{k}}{|\boldsymbol{N} \times \boldsymbol{k}|}\,. \tag{A1b}$$

Using $\boldsymbol{k}$, expressed in terms of $(\boldsymbol{e}_X, \boldsymbol{e}_Y, \boldsymbol{e}_Z)$ and $(\boldsymbol{p}, \boldsymbol{q}, \boldsymbol{N})$, given by

$$\begin{aligned}\boldsymbol{k} &= S_\Theta \sin\Upsilon \boldsymbol{e}_X - S_\Theta \cos\Upsilon \boldsymbol{e}_Y + C_\Theta \boldsymbol{e}_Z \\ &= S_\Theta \sin\Upsilon \boldsymbol{p} + (-S_{i_0} C_\Theta - C_{i_0} S_\Theta \cos\Upsilon)\boldsymbol{q} \\ &\quad + (C_{i_0} C_\Theta - S_{i_0} S_\Theta \cos\Upsilon)\boldsymbol{N}\,,\end{aligned} \tag{A2}$$

we compute the following expressions for $\boldsymbol{i}$ and $\boldsymbol{i}_i$ as

$$\begin{aligned}\boldsymbol{i} &= \cos\Upsilon \boldsymbol{e}_X + \sin\Upsilon \boldsymbol{e}_Y \\ &= \cos\Upsilon \boldsymbol{p} + C_{i_0} \sin\Upsilon \boldsymbol{q} + S_{i_0} \sin\Upsilon \boldsymbol{N}\,,\end{aligned} \tag{A3a}$$

$$\boldsymbol{i}_i = \frac{1}{\sin i}\left[(S_{i_0}C_\Theta + C_{i_0}S_\Theta\cos\Upsilon)\boldsymbol{p} + S_\Theta\sin\Upsilon\,\boldsymbol{q}\right] , \tag{A3b}$$

where $\sin i = |\boldsymbol{N}\times\boldsymbol{k}|$. The time evolution for the (observational) orbital inclination i is simply given by

$$\begin{aligned}\cos i &= \boldsymbol{N}\cdot\boldsymbol{k} = C_{i_0}C_\Theta - S_{i_0}S_\Theta\cos\Upsilon , && \text{(A4a)}\\ \sin i &= |\boldsymbol{N}\times\boldsymbol{k}| \\ &= \left[(S_{i_0}C_\Theta + C_{i_0}S_\Theta\cos\Upsilon)^2 + S_\Theta^2\sin^2\Upsilon\right]^{1/2} \\ &= \left[1-(C_{i_0}C_\Theta - S_{i_0}S_\Theta\cos\Upsilon)^2\right]^{1/2} . && \text{(A4b)}\end{aligned}$$

The equation for $\cos i$ is in agreement with Eq. (6) in Ref. [26]. We may also obtain a differential equation for the time evolution of i, as given in Ref. [10]. To get that, we differentiate $\cos i = \boldsymbol{N}\cdot\boldsymbol{k}$ with respect to time and make use of $\dot{\boldsymbol{k}} = \dot{\boldsymbol{L}}/L$ with $\dot{\boldsymbol{L}} = \boldsymbol{S}_{\rm eff}\times\boldsymbol{L}/(c^2r^3)$, $\sin i = |\boldsymbol{N}\times\boldsymbol{k}|$ and Eq. (A1b). In this way we deduce the following differential equation for the orbital inclination i:

$$\frac{di}{dt} = \frac{1}{c^2r^3}\boldsymbol{S}_{\rm eff}\cdot\frac{\boldsymbol{N}\times\boldsymbol{k}}{\sin i} = \frac{1}{c^2r^3}\boldsymbol{S}_{\rm eff}\cdot\boldsymbol{i}_i , \tag{A5}$$

which agrees with Eq. (5.15) of Ref. [10], whereas we note that the quantity characterizing the *ascending node* in Ref. [10], namely $\boldsymbol{i}$, is identical to our $\boldsymbol{i}_i$.

Following Ref. [10], let Ω be the longitude of the line of nodes $\boldsymbol{i}_i$, as measured from $\boldsymbol{p}$ in the $\boldsymbol{p}$-$\boldsymbol{q}$ plane (plane of the sky), which implies

$$\boldsymbol{i}_i = \cos\Omega\,\boldsymbol{p} + \sin\Omega\,\boldsymbol{q} . \tag{A6}$$

The time evolution for Ω is defined by either of the following two equations

$$\begin{aligned}\cos\Omega &= \boldsymbol{p}\cdot\boldsymbol{i}_i = \frac{1}{\sin i}(S_{i_0}C_\Theta + C_{i_0}S_\Theta\cos\Upsilon) , && \text{(A7a)}\\ \sin\Omega &= |\boldsymbol{p}\times\boldsymbol{i}_i| = \frac{1}{\sin i}S_\Theta\sin\Upsilon . && \text{(A7b)}\end{aligned}$$

Finally, we note that it is also possible to obtain the time evolution for the longitude of periastron ω, measured from $\boldsymbol{i}_i$ in the orbital plane, as defined in Ref. [10]. To do that, it is advantageous to define a periastron-based orthonormal triad $(\boldsymbol{a},\boldsymbol{b},\boldsymbol{k}) = (\boldsymbol{a},\boldsymbol{k}\times\boldsymbol{a},\boldsymbol{k})$. This is easily obtained by a rotation of the $(\boldsymbol{i},\boldsymbol{j},\boldsymbol{k})$ frame around the orbital angular momentum unit vector $\boldsymbol{k}$, by an angle Π. This gives us

$$\begin{pmatrix}\boldsymbol{a}\\ \boldsymbol{b}\\ \boldsymbol{k}\end{pmatrix} = \begin{pmatrix}\cos\Pi & \sin\Pi & 0\\ -\sin\Pi & \cos\Pi & 0\\ 0 & 0 & 1\end{pmatrix}\begin{pmatrix}\boldsymbol{i}\\ \boldsymbol{j}\\ \boldsymbol{k}\end{pmatrix} , \tag{A8}$$

where Π is the angle between $\boldsymbol{i}$ and the (instantaneous) periastron unit vector $\boldsymbol{a}$. The time evolution of ω is given by either of the following relations

$$\begin{aligned}\cos\omega &= \boldsymbol{i}_i\cdot\boldsymbol{a} = \frac{1}{\sin i}\left[-S_{i_0}\sin\Upsilon\sin\Pi\right.\\ &\quad\left.+(C_{i_0}S_\Theta + S_{i_0}C_\Theta\cos\Upsilon)\cos\Pi\right] , && \text{(A9a)}\\ \sin\omega &= |\boldsymbol{i}_i\times\boldsymbol{a}| = \frac{1}{\sin i}\left[(C_{i_0}S_\Theta + S_{i_0}C_\Theta\cos\Upsilon)\sin\Pi\right.\\ &\quad\left.+S_{i_0}\sin\Upsilon\cos\Pi\right] . && \text{(A9b)}\end{aligned}$$

The parametric solution to Π, which also gives its time evolution, is given by our parametrization via Eq. (3.1). In addition, the parametric solution to Π can be also obtained by noting that $\boldsymbol{a} = \boldsymbol{\mathcal{A}}/|\boldsymbol{\mathcal{A}}|$, where $\boldsymbol{\mathcal{A}}$ is the Laplace-Runge-Lenz vector given by

$$\begin{aligned}\boldsymbol{\mathcal{A}} &= \boldsymbol{\mathcal{P}}\times(\boldsymbol{\mathcal{R}}\times\boldsymbol{\mathcal{P}}) - \mu^2GM\frac{\boldsymbol{\mathcal{R}}}{\mathcal{R}}\\ &= \mu^2GM\left[\boldsymbol{p}\times(\boldsymbol{r}\times\boldsymbol{p}) - \frac{\boldsymbol{r}}{r}\right] . && \text{(A10)}\end{aligned}$$

Further details are described in Appendix B.

Eq. (A9b) for $\sin\omega$ is in agreement with Eq. (6) in Ref. [26]. The additional check $\sin^2\omega + \cos^2\omega = 1$ leads to

$$\sin i = \left[(C_{i_0}S_\Theta + S_{i_0}C_\Theta\cos\Upsilon)^2 + S_{i_0}^2\sin^2\Upsilon\right]^{1/2} , \tag{A11}$$

which is equivalent to Eq. (A4b) (using trigonometric relations).

Since $\boldsymbol{i}_i$, $\boldsymbol{i}$, and $\boldsymbol{a}$ lie in the orbital plane, we may define Δ such that it is the angle between $\boldsymbol{i}_i$ and $\boldsymbol{i}$. This implies that $\omega = \Pi + \Delta$ (see Fig. 1) and the time evolution for Δ is given by

$$\begin{aligned}\cos\Delta &= \boldsymbol{i}_i\cdot\boldsymbol{i} = \frac{1}{\sin i}(C_{i_0}S_\Theta + S_{i_0}C_\Theta\cos\Upsilon) , && \text{(A12a)}\\ \sin\Delta &= |\boldsymbol{i}_i\times\boldsymbol{i}| = \frac{1}{\sin i}S_{i_0}\sin\Upsilon . && \text{(A12b)}\end{aligned}$$

Finally, let $\varphi_{\rm DS}$ be the polar angle of $\boldsymbol{r}$, measured from $\boldsymbol{i}_i$, as defined by Damour and Schäfer in Ref. [10]. As we define φ from our $\boldsymbol{i}$, this leads to $\varphi_{\rm DS} = \varphi + \Delta$. Using these inputs, it is possible to write the relative separation vector $\boldsymbol{r}$ in $(\boldsymbol{a},\boldsymbol{b})$ in terms of $\varphi_{\rm DS}$ and ω as

$$\begin{aligned}\boldsymbol{r} &= r\cos(\varphi_{\rm DS}-\omega)\boldsymbol{a} + r\sin(\varphi_{\rm DS}-\omega)\boldsymbol{b} ,\\ &= r\cos v\,\boldsymbol{a} + r\sin v\,\boldsymbol{b} , && \text{(A13)}\end{aligned}$$

This is consistent with Eqs. (5.5b), (5.6), and (5.10) in Ref. [10]. In our angular variables, Eq. (A13) reads

$$\begin{aligned}\boldsymbol{r} &= r\cos(\varphi-\Pi)\boldsymbol{a} + r\sin(\varphi-\Pi)\boldsymbol{b} ,\\ &= r\cos v\,\boldsymbol{a} + r\sin v\,\boldsymbol{b} . && \text{(A14)}\end{aligned}$$

Appendix B: THE COMPUTATION OF $\dot{\Upsilon}$ AND $\dot{\Pi}$ VIA THE LAPLACE-RUNGE-LENZ VECTOR

In this section, we present an elegant way, starting from the famous Laplace-Runge-Lenz vector, to obtain the time evolution of the apsidal motion without any parametrization. The quantities in this section are non-reduced for a simple comparism with the work of Damour and Schäfer [10], and the work of Wex and Kopeikin [27].

We define the following functions of the canonically conjugate phase space variables $\boldsymbol{\mathcal{R}}$ and $\boldsymbol{\mathcal{P}}$

$$\boldsymbol{\mathcal{L}}(\boldsymbol{\mathcal{R}},\boldsymbol{\mathcal{P}}) \equiv \boldsymbol{\mathcal{R}} \times \boldsymbol{\mathcal{P}}\,, \tag{B1a}$$

$$\boldsymbol{\mathcal{A}}(\boldsymbol{\mathcal{R}},\boldsymbol{\mathcal{P}}) \equiv \boldsymbol{\mathcal{P}} \times (\boldsymbol{\mathcal{R}} \times \boldsymbol{\mathcal{P}}) - \mu^2 G M \frac{\boldsymbol{\mathcal{R}}}{\mathcal{R}}\,. \tag{B1b}$$

The first equation gives the orbital angular momentum vector $\boldsymbol{\mathcal{L}}$ and the second one the apsidal vector $\boldsymbol{\mathcal{A}}$, which is the famous supplementary first integral of the Kepler problem. The vector $\boldsymbol{\mathcal{A}}$ is usually named after Runge and Lenz or Laplace, though he was first discovered by Lagrange.

These vectors evolve according to the fundamental equations of Hamiltonian dynamics

$$\frac{d\boldsymbol{\mathcal{L}}}{dT} = \{\boldsymbol{\mathcal{L}},\mathcal{H}\}\,, \tag{B2a}$$

$$\frac{d\boldsymbol{\mathcal{A}}}{dT} = \{\boldsymbol{\mathcal{A}},\mathcal{H}\}\,, \tag{B2b}$$

where $\{\ldots,\ldots\}$ denotes the Poisson brackets and $\mathcal{H}$ is the full Hamiltonian for the relative motion of our binary system. As $\boldsymbol{\mathcal{L}}$ and $\boldsymbol{\mathcal{A}}$ are first integrals of $\mathcal{H}_{\rm N}$, only $\mathcal{H}_{\rm 1PN}+\mathcal{H}_{\rm 2PN}+\mathcal{H}_{\rm 3PN}+\mathcal{H}_{\rm SO}$ contribute to the right-hand sides of Eqs. (B2). The effects of $\mathcal{H}_{\rm 1PN}+\mathcal{H}_{\rm 2PN}+\mathcal{H}_{\rm 3PN}$ are already studied in the literature, and so we just want to consider the contribution of

$$\mathcal{H}_{\rm SO}(\boldsymbol{\mathcal{R}},\boldsymbol{\mathcal{P}},\boldsymbol{\mathcal{S}}_1,\boldsymbol{\mathcal{S}}_2) = \frac{G}{c^2\mathcal{R}^3}\boldsymbol{\mathcal{L}}\cdot\boldsymbol{\mathcal{S}}_{\rm eff}\,, \tag{B3}$$

where the effective spin $\boldsymbol{\mathcal{S}}_{\rm eff}$ is given by

$$\boldsymbol{\mathcal{S}}_{\rm eff} \equiv \left(2+\frac{3m_2}{2m_1}\right)\boldsymbol{\mathcal{S}}_1 + \left(2+\frac{3m_1}{2m_2}\right)\boldsymbol{\mathcal{S}}_2\,. \tag{B4}$$

We recall that $\boldsymbol{\mathcal{R}}$, $\boldsymbol{\mathcal{P}}$, $\boldsymbol{\mathcal{S}}_1$, and $\boldsymbol{\mathcal{S}}_2$ are canonical variables, such that the orbital variables commute with the spin variables, e.g., see Refs. [6, 22]. Using the basic Poisson brackets $\{\mathcal{R}^i,\mathcal{P}_j\} = \delta^i_j$, $\{\mathcal{S}^i_1,\mathcal{S}^j_1\} = \varepsilon^{ijk}\mathcal{S}^k_1$, $\{\mathcal{S}^i_2,\mathcal{S}^j_2\} = \varepsilon^{ijk}\mathcal{S}^k_2$, $\{\mathcal{R}^i,\mathcal{R}^j\} = \{\mathcal{P}_i,\mathcal{P}_j\} = \{\mathcal{S}^i_1,\mathcal{S}^j_2\} = \{\mathcal{R}^i,\mathcal{S}^j_1\} = \{\mathcal{R}^i,\mathcal{S}^j_2\} = \{\mathcal{P}_i,\mathcal{S}^j_1\} = \{\mathcal{P}_i,\mathcal{S}^j_2\} = 0$, we find

$$\{\boldsymbol{\mathcal{L}},\mathcal{H}_{\rm SO}\} = \frac{G}{c^2\mathcal{R}^3}\boldsymbol{\mathcal{S}}_{\rm eff}\times\boldsymbol{\mathcal{L}}\,, \tag{B5a}$$

$$\{\boldsymbol{\mathcal{A}},\mathcal{H}_{\rm SO}\} = \frac{G}{c^2\mathcal{R}^3}\left[\boldsymbol{\mathcal{S}}_{\rm eff}\times\boldsymbol{\mathcal{A}} - 3(\boldsymbol{\mathcal{L}}\cdot\boldsymbol{\mathcal{S}}_{\rm eff})\boldsymbol{\mathcal{L}}\times\frac{\boldsymbol{n}}{\mathcal{R}}\right]. \tag{B5b}$$

After orbital averaging these expressions, we can write down the equations for the precessional motion, induced by the spin-orbit interaction, in the vectorial form with the help of the rotational velocity vector $\boldsymbol{\Omega}_{\rm so}$

$$\left\langle\frac{d\boldsymbol{\mathcal{L}}}{dT}\right\rangle^{\rm so}_T = \boldsymbol{\Omega}_{\rm so}\times\boldsymbol{\mathcal{L}}\,, \tag{B6a}$$

$$\left\langle\frac{d\boldsymbol{\mathcal{A}}}{dT}\right\rangle^{\rm so}_T = \boldsymbol{\Omega}_{\rm so}\times\boldsymbol{\mathcal{A}}\,, \tag{B6b}$$

where

$$\boldsymbol{\Omega}_{\rm so} = \frac{G}{c^2}\left\langle\frac{1}{\mathcal{R}^3}\right\rangle_T\left[\boldsymbol{\mathcal{S}}_{\rm eff} - 3\frac{\boldsymbol{\mathcal{L}}\cdot\boldsymbol{\mathcal{S}}_{\rm eff}}{\mathcal{L}^2}\boldsymbol{\mathcal{L}}\right], \tag{B7}$$

and $\langle\ldots\rangle_T$ denotes the orbital average.

In case of the so-called *simple precession*, where the orbital angular momentum $\boldsymbol{\mathcal{L}}$ and the total spin $\boldsymbol{\mathcal{S}}$ precess about the conserved total angular momentum vector $\boldsymbol{\mathcal{J}} = \boldsymbol{\mathcal{L}}+\boldsymbol{\mathcal{S}}$ at the same rate [14, 23, 27], the precession of the orbital plane and the longitude of periastron are best described by the angles Υ and Π, defined with respect to the invariable plane, i.e., the plane perpendicular to $\boldsymbol{\mathcal{J}} = \mathcal{J}\boldsymbol{e}_Z$ (see Fig. 1 for details):

$$\frac{d\boldsymbol{\mathcal{L}}}{dT} = \frac{d\Upsilon}{dT}\boldsymbol{e}_Z\times\boldsymbol{\mathcal{L}}\,, \tag{B8a}$$

$$\frac{d\boldsymbol{\mathcal{A}}}{dT} = \left(\frac{d\Upsilon}{dT}\boldsymbol{e}_Z + \frac{d\Pi}{dT}\boldsymbol{k}\right)\times\boldsymbol{\mathcal{A}}, \tag{B8b}$$

where $\boldsymbol{k} = \boldsymbol{\mathcal{L}}/\mathcal{L}$ denotes the unit vector in the direction of the orbital angular momentum vector $\boldsymbol{\mathcal{L}}$. The conditions for the *simple precession* are fulfilled in our restricted cases, namely (i), the equal-mass case and (ii), the single-spin case. In these cases the effective spin $\boldsymbol{\mathcal{S}}_{\rm eff}$ and the total spin $\boldsymbol{\mathcal{S}}$ are related by $\boldsymbol{\mathcal{S}}_{\rm eff} = \lambda_{\rm so}\boldsymbol{\mathcal{S}}$, where the mass dependent coupling constant is given by

$$\lambda_{\rm so} \equiv \begin{cases} 2+3/2 = 7/2 & \text{for (i), the equal-mass case}\,,\\ 2+3m_2/(2m_1) & \text{for (ii)}, \boldsymbol{\mathcal{S}}_1 \neq 0\,, \boldsymbol{\mathcal{S}}_2 = 0\,,\\ 2+3m_1/(2m_2) & \text{for (ii)}, \boldsymbol{\mathcal{S}}_1 = 0\,, \boldsymbol{\mathcal{S}}_2 \neq 0\,. \end{cases}$$

Note, that $\lambda_{\rm so} = \chi_{\rm so}/\eta$. Using $\boldsymbol{\mathcal{S}}_{\rm eff} = \lambda_{\rm so}\boldsymbol{\mathcal{S}}$ in Eq. (B7) and comparing Eqs. (B6) with an averaged version of Eqs. (B8), we obtain the precession in terms of the angles Υ and Π

$$\left\langle\frac{d\Upsilon}{dT}\right\rangle^{\rm so}_T = \frac{G}{c^2}\left\langle\frac{1}{\mathcal{R}^3}\right\rangle_T\lambda_{\rm so}\mathcal{J}\,, \tag{B9a}$$

$$\left\langle\frac{d\Pi}{dT}\right\rangle^{\rm so}_T = \frac{G}{c^2}\left\langle\frac{1}{\mathcal{R}^3}\right\rangle_T\lambda_{\rm so}\mathcal{L}\left(-1-3\frac{\boldsymbol{\mathcal{L}}\cdot\boldsymbol{\mathcal{S}}}{\mathcal{L}^2}\right). \tag{B9b}$$

We note, that in case of $d\Upsilon/dT$ it is even possible to obtain a non-averaged equation, if one uses Eq. (B5a) instead of Eq. (B6a) for the comparism with Eq. (B8a).

Our results, given by Eqs. (B9), are in agreement with Eqs. (4.11) and with Eqs. (38) and (42) in Ref. [27].

---

[1] I. H. Stairs, Living Rev. Relativ. **6**, 5 (2003).

[2] Following URLs provide a wealth of information about the ground and space based gravitational wave interferometers: `http://www.ligo.caltech.edu`, `http://www.virgo.infn.it`, `http://www.geo600.uni-hannover.de`, `http://tamago.mtk.nao.ac.jp`, and `http://lisa.nasa.gov`.

[3] M. Ansorg, Phys. Rev. D **72**, 024018 (2005).

[4] J. Levin, Phys. Rev. D **67**, 044013 (2003).

[5] A. Gopakumar and C. Königsdörffer, *The deterministic nature of conservative post-Newtonian accurate dynamics of compact binaries with leading order spin-orbit interaction*, submitted to Phys. Rev. D.

[6] For a brief introduction to spin effects in general relativity, see G. Schäfer, Gen. Relativ. Gravit. **36**, 2223 (2004), gr-qc/0407116.

[7] T. Damour and N. Deruelle, Ann. Inst. Henri Poincaré Phys. Théor. **43**, 107 (1985).

[8] T. Damour and N. Deruelle, Ann. Inst. Henri Poincaré Phys. Théor. **44**, 263 (1986).

[9] T. Damour and J. Taylor, Phys. Rev. D **45**, 1840 (1992).

[10] T. Damour and G. Schäfer, Nuovo Cimento Soc. Ital. Fis., B **101**, 127 (1988).

[11] G. Schäfer and N. Wex, Phys. Lett. A **174**, 196, (1993); **177**, 461(E) (1993).

[12] T. Damour, A. Gopakumar, and B. R. Iyer, Phys. Rev. D **70**, 064028 (2004).

[13] R.-M. Memmesheimer, A. Gopakumar, and G. Schäfer, Phys. Rev. D **70**, 104011 (2005).

[14] C. Königsdörffer and A. Gopakumar, Phys. Rev. D **71**, 024039 (2005).

[15] C. Königsdörffer and A. Gopakumar (to be published).

[16] J. M. Cordes, M. Kramer, T. J. W. Lazio, B. W. Stappers, D. C. Backer, and S. Johnston, New Astron. Rev. **48**, 1413 (2004).

[17] M. Burgay *et al*, Nature **426**, 531 (2003).

[18] A. G. Lyne *et al*, Science **303**, 1153 (2004).

[19] I. A. Morrison, T. W. Baumgarte, S. L. Shapiro, and V. R. Pandharipande, Astrophys. J. **617**, L135 (2004).

[20] J. M. Lattimer and B. F. Schutz, *Constraining the equation of state with moment of inertia measurements*, astro-ph/0411470.

[21] G. Schäfer, Ann. Phys. (N.Y.) **161**, 81 (1985).

[22] T. Damour, Phys. Rev. D **64**, 124013 (2001).

[23] T. A. Apostolatos, C. Cutler, G. J. Sussman, and K. S. Thorne, Phys. Rev. D **49**, 6274 (1994).

[24] N. Wex, Class. Quantum Grav. **12**, 983 (1995).

[25] H. Goldstein, *Classical Mechanics*, 2nd edition (Addison-Wesley, Reading, Mass., 1980).

[26] D. Lai, L. Bildsten, and V. M. Kaspi, Astrophys. J. **452**, 819 (1995).

[27] N. Wex and S. M. Kopeikin, Astrophys. J. **514**, 388 (1999).